

## Physician - Heart Failure, Transplantation and Mechanical Support Systems

[MSB-06]

### Retrospective Cohort Analysis of Ventricular Assist Device Driveline Infections: A Single-Center Experience

Melisa Kandemir<sup>1</sup>, Mehmet Cahit Sarıcaoğlu<sup>1</sup>, Mukime Elif Sarıcaoğlu<sup>2</sup>, Mustafa Bahadır İnan<sup>1</sup>, Tuğba Altuntaş<sup>3</sup>, Zeynep Eyileten<sup>1</sup>, Sadık Eryılmaz<sup>1</sup>, Ahmet Rüçhan Akar<sup>1</sup>

<sup>1</sup>Department of Cardiovascular Surgery, Ankara University School of Medicine, Ankara, Türkiye

<sup>2</sup>Department of Infectious Diseases and Clinical Microbiology, Ankara University School of Medicine, Ankara, Türkiye

<sup>3</sup>Organ Transplant Unit, Ankara University School of Medicine, Ankara, Türkiye

---

*Cardiovascular Surgery and Interventions* 2024;11(Suppl 1):MSB-06

Doi: 10.5606/e-cvsi.2024.msb-06

E-mail: melisakandemir1999@gmail.com

Received: August 04, 2024 - Accepted: September 29, 2024

---

**Objective:** This study aimed to discuss our approach to driveline infections (DLIs), which can be managed by debridement, open packing, delayed closure, closed catheter irrigation, and rectus muscle relocation, and define the risk factors.

**Methods:** This retrospective study reviewed all patients who underwent ventricular assist device (VAD) implantation at our institution between May 2011 and May 2023. Demographic data, comorbidities, VAD type, time from VAD implantation to infection, microbiological testing, surgical debridement, antibiotic therapy, infectious symptoms, hospital readmissions, postoperative complications, and overall survival were evaluated.

**Results:** During the study period, 90 patients underwent VAD implantation, and DLIs were detected in 20 patients. The mean VAD time was 561 days in all patients, and the mean VAD time was 1277 days in those with DLIs. A statistically significant relationship was found between the risk of DLI and the total duration of VAD time ( $p < 0.05$ ). Median time from VAD implantation to first DLI admission was 513 days. Most commonly detected pathogens were *Staphylococcus* spp. (64%) and *Pseudomonas aeruginosa* (25%). Surgical debridement was performed in 10 patients; nine had driveline relocation followed by vacuum-assisted closure until their culture results were negative.

**Conclusion:** Driveline infections endanger the final objectives of VAD therapy by disrupting patient autonomy, the chance of bridging to heart transplantation, and decreasing overall survival and quality of life. Management of DLI usually requires extended, repeated hospitalizations and intense outpatient care. Vacuum therapy and muscle relocation have emerged as essential adjuncts to treating DLIs. Surgical treatment modalities should be standardized.

**Keywords:** Driveline infections, surgical debridement, ventricular assist device.

#### References

1. Roussel LO, Khouri JS, Christiano JG. Pedicled flap closure as an adjunct for infected ventricular assist devices. *Ann Plast Surg* 2017;78:712-6. doi: 10.1097/SAP.0000000000000904.
2. Hieda M, Sata M, Nakatani T. The importance of the management of infectious complications for patients with left ventricular assist device. *Healthcare (Basel)* 2015;3:750-6. doi: 10.3390/healthcare3030750.
3. Leuck AM. Left ventricular assist device driveline infections: Recent advances and future goals. *J Thorac Dis* 2015;7:2151-7. doi: 10.3978/j.issn.2072-1439.2015.11.06.
4. Koval CE, Thuita L, Moazami N, Blackstone E. Evolution and impact of drive-line infection in a large cohort of continuous-flow ventricular assist device recipients. *J Heart Lung Transplant* 2014;33:1164-72. doi: 10.1016/j.healun.2014.05.011.
5. Parameshwar J, Hogg R, Rushton S, Taylor R, Shaw S, Mehew J, et al. Patient survival and therapeutic outcome in the UK bridge to transplant left ventricular assist device population. *Heart (British Cardiac Society)* 2019;105:291-296. doi: 10.1136/heartjnl-2018-313355.