

Physician - Valvular Diseases and Surgery

[MSB-25]

Mid-Term Results of Ozaki Procedure: Azerbaijan Experience

Kamran Ahmadov, Nigar Kazimzade, Kamran Musayev

Department of Cardiovascular Surgery, Merkezi Klinika, Baki, Azerbaijan

Cardiovascular Surgery and Interventions 2024;11(Suppl 1):MSB-25

Doi: 10.5606/e-cvsi.2024.msb-25

E-mail: Dr.kamran.ahmadov@gmail.com

Received: September 09, 2024 - Accepted: September 29, 2024

Objective: This study aimed to report the outcomes of the Ozaki procedure, which involves reconstructing aortic valve leaflets with autologous pericardium, performed in a clinic in Azerbaijan.

Methods: This retrospective study analyzed 40 patients who underwent aortic valve reconstruction between August 2018 and June 2023. Patients were divided into two groups: Group A (mean age: 63 years) followed the traditional Ozaki technique, while Group B (mean age: 65 years) received an additional commissural reinforcement, a modification proposed by our team.

Results: Presenting symptoms were aortic stenosis or a combination of aortic stenosis and aortic regurgitation. Preoperative echocardiography showed peak and mean pressure gradients of 84 ± 34.6 and 50.5 ± 23 mmHg, respectively. Cardiopulmonary bypass and aortic cross-clamp times averaged 142 and 115 min for Group A and 144 and 107 min for Group B. There were no in-hospital mortalities or pacemaker implantations. No significant increases in aortic gradients were noted, and no reoperations were required. Four patients in Group A developed mild aortic regurgitation during follow-up, while in Group B, aortic regurgitation remained minimal or absent. The median follow-up periods were 64 months for Group A and 28 months for Group B. The study demonstrated 100% freedom from major adverse valve-related events during the follow-up.

Conclusion: Since its introduction, aortic valve neocuspidization has gained popularity. Mid-term results from the Ozaki procedure showed favorable outcomes in terms of mortality, valve gradients, and freedom from adverse valve-related events. While some studies have reported a slight increase in valve regurgitation following the Ozaki procedure, our novel additional commissural reinforcement technique provided a reduction in aortic valve regurgitation during follow-up. Further studies are needed to assess long-term results.

Keywords: Aortic valve, aortic valve disease.




Figure 1. Image of the reconstructed valve with additional commissural reinforcement.

References

1. Ozaki S, Kawase I, Yamashita H, Uchida S, Takatoh M, Hagiwara S, et al. Aortic valve reconstruction using autologous pericardium for aortic stenosis. *Circ J* 2015;79:1504-10. doi: 10.1253/circj.CJ-14-1092.
2. Mokryk I, Nechai I, Stetsyuk I, Malova N, Demyanchuk V, Todurov B. Complete aortic valve reconstruction with autologous pericardium: Analysis of mid-term results of single-center experience with AVNeo procedure. *Ann Thorac Cardiovasc Surg* 2024;30:24-00067. doi: 10.5761/atcs.aa.24-00067.
3. Al Halees Z, Al Shahid M, Al Sanei A, Sallehuddin A, Duran C. Up to 16 years follow-up of aortic valve reconstruction with pericardium: A stentless readily available cheap valve? *Eur J Cardiothorac Surg*. 2005;28:200-5. doi: 10.1016/j.ejcts.2005.04.041.