

## Physician - Peripheral Artery Diseases and Surgery/Percutan Interventions

[MEP-47]

### Anterior Prosthetic Knee Dislocation with Popliteal Artery Injury: A Case Report

Ali Fuat Karaçuha<sup>1</sup>, Mehmet Cahit Sarıcaoğlu<sup>2</sup>, Evren Özçınar<sup>2</sup>, Levent Yazıcıoğlu<sup>2</sup>, Sadık Eryılmaz<sup>2</sup>

<sup>1</sup>Department of Cardiovascular Surgery, Trabzon Kanuni Training and Research Hospital, Trabzon, Türkiye

<sup>2</sup>Department of Cardiovascular Surgery, Ankara University Faculty of Medicine, Ankara, Türkiye

*Cardiovascular Surgery and Interventions* 2024;11(Suppl 1):MEP-47

Doi: 10.5606/e-cvsi.2024.mep-47

E-mail: alifuatkaracuha@hotmail.com

Received: September 14, 2024 - Accepted: September 29, 2024

Anterior dislocation is very rare in patients with a knee prosthesis. Few cases of acute vascular injury after anterior prosthesis dislocation have been reported in the literature with extremely poor outcomes. Herein, we reported the case of a 69-year-old female who developed an anterior knee prosthesis dislocation after a slip and fall at home, injuring the popliteal artery, which was spontaneously reduced. Following orthopedic controls, the patient was taken to the hybrid operating room, and digital subtraction angiography showed a transection of the right popliteal artery with no distal flow. Emergency fasciotomy and popliteal bypass were performed with simultaneous external fixation of the joint. Distal flow was restored and the limb was saved. External fixation of the knee allowed better stabilization of the joint, and preservation of the repaired vascular injury. Although there are no clear guidelines for management, we emphasize the importance of the hybrid operating room for rapid diagnosis of vascular trauma and rapid transition to open surgery to repair the injury.

**Keywords:** Injury, knee prosthesis, popliteal artery, trauma.



**Figure 1.** Digital subtraction angiography.

## References

1. Makaloski V, Stellmes A, Wyss D, Weiss S, Becker D, Wyss TR, et al. Posterior approach for revascularization in blunt popliteal vessel injury. *Ann Vasc Surg* 2018;48:89-96. doi: 10.1016/j.avsg.2017.10.019.
2. Frykberg ER. Popliteal vascular injuries. *Surg Clin North Am* 2002;82:67-89. doi: 10.1016/S0039-6109(03)00141-5.
3. Sillanpää PJ, Kannus P, Niemi ST, Rolf C, Felländer-Tsai L, Mattila VM. Incidence of knee dislocation and concomitant vascular injury requiring surgery: A nationwide study. *J Trauma Acute Care Surg* 2014;76:715-9. doi: 10.1097/TA.0000000000000136.
4. Canovas F, Dagneaux L. Quality of life after total knee arthroplasty. *Orthop Traumatol Surg Res* 2018;104:S41-6. doi: 10.1016/j.otsr.2017.04.017.
5. Trickett JP, Scott RA, Tilney HS. Screening and management of asymptomatic popliteal aneurysms. *J Med Screen* 2002;9:92-3. doi: 10.1136/jms.9.2.92.
6. Brown SL, Lewis M, Morrow DR. Endovascular repair of ruptured popliteal artery aneurysms: A case report and review of the literature. *EJVES Short Rep* 2016;32:24-8. doi: 10.1016/j.ejvssr.2016.07.001.
7. Chapman JA. Popliteal artery damage in closed injuries of the knee. *J Bone Joint Surg Br* 1985;67:420-3. doi: 10.1302/0301-620X.67B3.3997952.
8. Ahn RS, Brown MJ, Santilli MD. Traumatic anterior knee dislocation after total knee arthroplasty. *Arthroplast Today* 2015;2:97-9. doi: 10.1016/j.artd.2015.10.006.
9. Nicandri GT, Chamberlain AM, Wahl CJ. Practical management of knee dislocations: A selective angiography protocol to detect limb-threatening vascular injuries. *Clin J Sport Med* 2009;19:125-9. doi: 10.1097/JSM.0b013e31819cd37a.