



Clues on electrocardiography to predict the presence of paroxysmal atrial fibrillation in patients with acute ischemic stroke: A propensity score-matched study

Oktay Şenöz , Ferhat Siyamend Yurdam 

Department of Cardiology, İzmir Bakırçay University, İzmir, Turkey

Received: January 21, 2022 Accepted: February 06, 2022 Published online: March 18, 2022

ABSTRACT

Objectives: In this study, we aimed to detect surface electrocardiography (ECG) markers that could be predictive of paroxysmal atrial fibrillation (PAF) attacks in patients with ischemic stroke.

Patients and methods: Between November 2017 and April 2021, a total of 112 patients (65 males, 47 females; mean age: 70.5±6.8 years; range, 51 to 84 years) hospitalized for acute ischemic stroke with sinus rhythm on surface ECG who underwent Holter ECG monitoring for PAF were retrospectively analyzed. The patients were divided into two groups of 56 patients in each (Group 1: those with PAF on Holter ECG and Group 2: those without PAF). Both groups were matched according to demographic, clinical, and echocardiographic features using the propensity score matching method.

Results: Demographic, clinical, and echocardiographic features were similar between groups ($p>0.05$). The mean maximum P-wave duration (PWD) and P-wave dispersion (PWD_{is}) were longer in Group 1 than Group 2 (108.4±9.9 *vs.* 102.5±10.2 ms; $p=0.002$, 49.4±13.6 *vs.* 36.8±11.7 ms; $p<0.001$). Similarly, the mean P-wave terminal force in lead V₁ (PTFV₁) was higher in Group 1 than Group 2 (4415±909 *vs.* 3826±568 $\mu V \cdot ms$; $p<0.001$). Logistic regression analysis revealed high PWD_{is} (odds ratio [OR]: 1.164; 95% confidence interval [CI]: 1.069-1.268; $p<0.001$) and PTFV₁ (OR: 1.156; 95% CI: 1.065-1.254; $p=0.001$) as independent predictors of PAF.

Conclusion: PWD_{is} and PTFV₁ are independent predictors of PAF in patients with acute ischemic stroke. These simple and easily accessible predictors that can be detected via surface ECG may be used as a guide to identify patients who require longer rhythm monitoring to better detect occult PAF, thereby preventing recurrent strokes.

Keywords: Atrial fibrillation, ischemic stroke, P-wave dispersion, P-wave terminal force in lead V₁.

Atrial fibrillation (AF) is one of the most frequent cardiac rhythm abnormalities. Although its prevalence in the general population is thought to be around 2 to 4%, it is expected that it would increase by 2.3 times as life expectancy increases and advanced investigation techniques lead to overdiagnosis.^[1] Diabetes mellitus, advanced age, hypertension, chronic renal failure, coronary artery disease, heart failure, obesity and obstructive sleep apnea are all important predisposing factors for AF.^[1,2] Also, AF can lead to serious complications such as heart failure, stroke and death.^[3] While the risk of AF associated stroke is 1.5% in the sixth decade of life, this risk increases to 24% in the ninth decade.^[4] Similar to clinically overt AF, paroxysmal atrial fibrillation (PAF) attacks also increase the risk of ischemic stroke. Current guidelines recommend monitoring with Holter electrocardiography (ECG) to detect PAF attacks in patients with ischemic stroke, even if the surface ECG is normal.^[5] Currently,

24 to 48-h Holter ECG monitoring is used for this purpose. However, existing silent PAF attacks cannot be detected with these recordings in some cases. Identifying patients at high risk for the development of AF before resorting to Holter ECG monitoring may help to improve diagnostic accuracy. In such high-risk patients, monitoring may be extended, if AF is not detected on 24 to 48-h Holter ECG recordings, thereby preventing false-negative results.

Many previous studies have investigated the use of several different scoring systems to identify patients

Corresponding author: Oktay Şenöz, MD. İzmir Bakırçay Üniversitesi, Kardioloji Anabilim Dalı, 35665 Menemen, İzmir, Türkiye.
Tel: +90 506 - 882 81 29 e-mail: oktayssss@hotmail.com

Citation:

Şenöz O, Yurdam FS. Clues on electrocardiography to predict the presence of paroxysmal atrial fibrillation in patients with acute ischemic stroke: A propensity score-matched study. *Cardiovasc Surg Int* 2022;9(1):36-42.

who are under high risk for development of AF.^[6,7] However, many of these systems are complex and difficult to use in daily practice. In the present study, we aimed to determine surface ECG findings that could predict the presence of PAF in those with ischemic stroke.

PATIENTS AND METHODS

This single-centre, retrospective study was conducted at Izmir Bakırçay University, Department of Cardiology between November 2017 and April 2021. A total of 149 patients with an ischemic cerebrovascular event (CVE) that had sinus rhythm on baseline surface ECG and underwent 24 to 48-h Holter ECG for cardioembolic investigation were screened. The patients were matched with the propensity score matching method in terms of their baseline clinical, demographic, and echocardiographic characteristics, which may be risk factors for AF, at a ratio of 1:1, and 56 each with and without PAF on Holter ECG a total of 112 patients (65 males, 47 females; mean age: 70.5 ± 6.8 years; range, 51 to 84 years) were included in the study. Those with PAF detected on Holter ECG were defined as Group 1 ($n=56$) and those without it were defined as Group 2 ($n=56$). Patients under 18 years of age, those with serious renal or hepatic insufficiency, rheumatic moderate to severe mitral valve stenosis, prosthetic heart valve, previous history of AF, those who did not have normal sinus rhythm on basal ECG and those with missing or insufficient data on ECG or Holter ECG records were excluded from the study. The study protocol was approved by the Bakırçay University Non-Interventional Clinical Research Ethics Committee (No: 2021-333). The study was conducted in accordance with the principles of the Declaration of Helsinki.

Patients' clinical, demographic, laboratory and ECG were obtained from the patient records. Baseline 12-lead surface ECG records were analyzed in detail. The 24 to 48-h Holter ECG recordings were evaluated by two cardiology specialists with the presence and length of PAF attacks noted.

Electrocardiography

During hospitalization with patients in the supine position, a 12-lead ECG was recorded from all subjects using 10 mV/cm standardization with 25 mm/sec paper speed and 0.05 to 100 Hz filter band settings. The ECGs were scanned and transferred to the

computer system and measurements were made with an electronic caliper at 4× magnification on the high-resolution computer screen. P-wave dispersion (PWD_{is}) was calculated as the difference between the maximum and minimum P wave duration (PWD) in all 12 ECG leads. P-wave terminal force in lead V₁ (PTFV₁) was defined as the duration multiplied by the amplitude of the negative part of the P wave (P'duration (a) × P'amplitude (b)), measured in μV ms in lead V₁ (Figure 1).

Detection of an AF attack of at least 30 sec on Holter ECG was defined as PAF.

Statistical analysis

Statistical analysis was performed using the IBM SPSS for Windows version 25.0 software (IBM Corp., Armonk, NY, USA). A propensity score for the presence of PAF was estimated for each patient with logistic regression, using 26 clinically relevant baseline variables. Thereafter, using 1:1 matching without replacement, a matched cohort was constructed matching each patient without PAF to the closest patient with PAF in which propensity score differed by 0.1 or less. The ability to balance baseline features was evaluated by absolute standardized differences (the percentage difference between the means for the two groups divided by the reciprocal standard deviation). In standard differences, 10% were considered inconsequential. After matching, the overall balance p value was calculated as 0.99.

The normality distribution of continuous variables was evaluated using the Kolmogorov-Smirnov test. Continuous variables were expressed in mean \pm standard deviation (SD) and categorical variables were expressed in number and frequency. The groups were compared using the independent Student t-test or Mann-Whitney U test according to the normality distribution for continuous variables, and the chi-square test or Fisher exact test for categorical



Figure 1. P-wave terminal force measurement in lead V₁ (PTFV₁) = (a) P'duration × (b) P'amplitude.

PTFV₁: P wave terminal force in lead V₁.

variables. The ECG findings were evaluated with logistic regression analysis to identify independent predictors of the presence of PAF on Holter ECG. Optimal cut-off values were determined by receiving operating characteristics (ROC) curve analysis to predict PAF. A *p* value of <0.05 was considered statistically significant.

RESULTS

Previous to ischemic stroke, 61.6% of patients (n=69) had a diagnosis of hypertension, 41.1% (n=46) had hyperlipidemia, and 33.9% (n=38) had diabetes mellitus. A previous history of CVE was present in 17.9% (n=20) patients. Coronary artery disease was present in 24.1% (n=27) and chronic heart failure in

Table 1
Baseline clinical characteristics and medications of matched patients

Variables	Group 1 (n=56)			Group 2 (n=56)			Standard differences	p
	n	%	Mean±SD	n	%	Mean±SD		
Clinical features								
Age (year)			70.6±7.1			70.2±6.5	0.052	0.782
Sex								
Male	33	58.9		32	57.1		0.017	0.848
Hypertension	33	58.9		36	64.3		0.053	0.560
Diabetes mellitus	19	33.9		19	33.9		0	1
Hypercholesterolemia	24	42.9		22	39.3		0.035	0.701
Chronic renal failure	8	14.3		9	16.1		0.017	0.792
CVD history	9	16.1		11	19.6		0.035	0.622
Coronary artery disease	14	25		13	23.2		0.017	0.825
Chronic heart failure	4	7.1		3	5.4		0.017	0.696
Thyroid disease	2	3.6		2	3.6		0	1
COPD	1	1.8		1	1.8		0	1
Carotid artery disease	26	46.4		28	50		0.071	0.611
CHA2DS2-VASc score			4.9±1.2			4.8±1.3	0.013	0.826
LVEF (%)			57.7±5.7			58.1±4.2	0.048	0.779
Left atrium diameter (mm)			37.1±3.5			37.1±4.4	0.004	0.981
LVDD	28	50		27	48.2		0.017	0.850
Severe mitral regurgitation	1	1.8		1	1.8		0	1
Medications								
Acetylsalicylic acid	13	23.2		15	26.8		0.035	0.663
Clopidogrel	10	17.9		9	16.1		0.017	0.801
Oral anticoagulant	2	3.6		1	1.8		0.017	0.390
Statin	11	19.6		9	16.1		0.035	0.622
Beta-blocker	13	23.2		14	25		0.017	0.825
Calcium channel blocker	19	33.9		18	32.1		0.017	0.841
ACEi or ARB	18	32.1		21	37.5		0.053	0.552
Levothyroxine	1	1.8		1	1.8		0	1
Methimazole	1	1.8		0	0		0.017	0.315
SD: Standard deviation; CVD: Cerebrovascular disease; COPD: Chronic obstructive pulmonary disease; LVEF: Left ventricular ejection fraction; LVDD: Left ventricular diastolic dysfunction; ACEi: Angiotensin converting enzyme inhibitor; ARB: Angiotensin receptor blocker.								

SD: Standard deviation; CVD: Cerebrovascular disease; COPD: Chronic obstructive pulmonary disease; LVEF: Left ventricular ejection fraction; LVDD: Left ventricular diastolic dysfunction; ACEi: Angiotensin converting enzyme inhibitor; ARB: Angiotensin receptor blocker.

Variables	Group 1 (n=56)	Group 2 (n=56)	<i>p</i>
	Mean±SD	Mean±SD	
Fasting blood glucose (mg/dL)	125.1±37.4	135.5±49.2	0.214
Urea (mg/dL)	39.8±17.5	42.9±23.2	0.370
Creatinine (mg/dL)	1.1±0.8	0.9±0.3	0.207
Sodium (mEq/L)	138.9±2.9	138.6±3.0	0.653
Potassium (mg/dL)	4.4±0.6	4.3±0.4	0.166
Total cholesterol (mg/dL)	183.0±40.1	195.5±41.0	0.822
LDL-cholesterol (mg/dL)	111.9±34.7	116.5±35.4	0.723
Plasma triglycerides (mg/dL)	143.5±59.5	193.2±101.1	0.009
HDL-cholesterol (mg/dL)	41.7±11.2	39.1±9.4	0.198
TSH (mU/L)	2.2±4.8	1.9±1.6	0.715
Free T4 (ng/dL)	1.3±0.4	1.2±0.2	0.961
White blood cell count (×10 ⁹ /L)	9.0±2.4	8.4±2.9	0.497
Neutrophil count (×10 ⁹ /L)	6.0±2.1	5.5±2.6	0.558
Lymphocyte count (×10 ⁹ /L)	2.0±0.8	2.0±0.9	0.938
Hemoglobin (g/dL)	12.9±1.6	13.0±1.9	0.224
Hematocrit (%)	38.4±4.5	38.6±5.3	0.199
Platelet count (×10 ⁹ /L)	266.1±90.7	253.6±84.8	0.369
SD: Standard deviation; LDL: Low-density lipoprotein; HDL: High-density lipoprotein; TSH: Thyroid-stimulating hormone.			

6.3% (n=7) patients. Fifty-four (48.2%) patients had carotid artery disease and, of these patients, 34 had carotid artery stenosis <50% and 20 had stenosis ≥50%. The mean CHA2DS2-VASc score of patients was 4.9±1.3. The mean left ventricular ejection fraction was 57.9±5.1% and the mean left atrial (LA) diameter was 37.1±4.1 mm. When CVE occurred, 25% (n=28) patients were using acetylsalicylic acid, 17% (n=19) clopidogrel and 2.7% (n=3) were taking

oral anticoagulant therapy. Patients' demographic and clinical features are shown in Table 1 and their laboratory findings are shown in Table 2.

Both groups were similar with regard to the mean baseline heart rate (74.4±13.2 *vs.* 74.6±10.8 bpm, *p*=0.93). The mean maximum PWD and PWD_{is} were longer in Group 1, compared to Group 2 (108.4±9.9 *vs.* 102.5±10.2 ms; *p*=0.002 and 49.4±13.6 *vs.* 36.8±11.7 ms; *p*<0.001, respectively). Similarly,

Variables	Group 1 (n=56)	Group 2 (n=56)	<i>p</i>
	Mean±SD	Mean±SD	
Heart rate (bpm)	74.4±13.2	74.6±10.8	0.931
Maximum P wave duration (ms)	108.4±9.9	102.5±10.2	0.002
P wave dispersion (ms)	49.4±13.6	36.8±11.7	<0.001
PTFV ₁ (µV·ms)	4415±909	3826±568	<0.001
SD: Standard deviation; bpm: Beats per minute; ms: Millisecond; PTFV ₁ : P wave terminal force in lead V ₁ .			

Table 4
Effects of ECG indicators on the detection of paroxysmal atrial fibrillation in univariate and multivariate logistic regression analysis

Variables	Univariate logistic regression			Multivariate logistic regression		
	OR	95% CI	<i>p</i>	OR	95% CI	<i>p</i>
Maximum P-wave duration	1.061	1.019-1.104	0.004	0.913	0.825-1.010	0.078
P-wave dispersion	1.082	1.044-1.121	<0.001	1.164	1.069-1.268	<0.001
PTFV ₁	1.109	1.042-1.180	0.001	1.156	1.065-1.254	0.001

OR: Odds ratio; CI: Confidence interval; PTFV₁: P wave terminal force in lead V₁.

PTFV₁ was higher in Group 1, compared to Group 2 (4415±909 *vs.* 3826±568 $\mu\text{V}\cdot\text{ms}$; $p<0.001$) (Table 3).

Logistic regression analysis demonstrated high PWD_{is} (odds ratio [OR]: 1.164; 95% confidence interval [CI]: 1.069-1.268; $p<0.001$) and high PTFV₁ (OR: 1.156; 95% CI: 1.065-1.254; $p=0.001$) as being independent predictors for the presence of PAF (Table 4). As predictors for the presence of PAF, the ROC curve analysis revealed a sensitivity of 71% and specificity of 69% for PWD_{is} >42 ms (area under the curve [AUC] 0.764, 95% CI: 0.675-0.852, $p<0.001$) and a sensitivity of 60% and specificity of 64%

for PTFV₁ >4.050 $\mu\text{V}\cdot\text{ms}$ (AUC: 0.686, 95% CI: 0.574-0.798, $p=0.002$) (Figure 2).

DISCUSSION

In the present study, we found that PWD_{is} and PTFV₁ parameters as measured on surface ECG were independent predictors for the presence of PAF in patients with ischemic stroke.

Some patients with AF describe palpitations, shortness of breath and fatigue while some may be completely asymptomatic and present with complications such as ischemic stroke or tachycardiomyopathy.^[8] Non-valvular AF is responsible for about half of all cardioembolic events.^[9] The incidence of occult or subclinical AF is not known. Therefore, patients with symptomatic AF that are observed in daily practice may be considered the tip of the iceberg. The development of new devices and applications has led to an increase in diagnosis rates of asymptomatic and subclinical AF. The rates of subclinical AF was reported to be 35% in a group of patients with implanted cardiac devices that were followed for 2.5 years.^[10] In patients with cryptogenic stroke, 12.5% were found to have PAF attacks during one-year rhythm monitorization.^[11] It is important to identify patients who do not have arrhythmia on surface ECG, but who are at high risk for the development of AF and to perform long-term rhythm monitoring in these patients to prevent ischemic stroke.

Many scoring methods have been utilized to predict the development of AF in those with normal surface ECG.^[6,7] The CHADS₂ and the CHA₂DS₂-VASc risk scores have been reported for prediction of new occurrence of AF, ischemic stroke and long-term outcomes after AF ablation.^[12] Christophersen et




Figure 2. Receiver operating characteristic (ROC) curve analysis of the optimal cut-off values of P-wave dispersion and P-wave terminal force in lead V₁.

PWD_{is}: P-wave dispersion (PWD_{is}; PTFV₁: P wave terminal force in lead V₁).

al.^[12] reported that CHARGE-AF scoring was better at predicting AF, compared to CHA2DS2-VASc. On the other hand, some studies have used the HATCH score for prediction of AF recurrence and persistence.^[13] The main feature of these scoring methods is that they predict the development of AF according to the clinical characteristics of the patients. However, AF is an ECG disorder and using ECG findings for its prediction may be a more plausible way. Electrocardiographic evaluation is also a simpler, cheaper, and easily accessible method than the aforementioned scoring systems. Furthermore, it has been reported that P-wave indices are as effective as clinical scoring methods for the prediction of AF and ischemic stroke.^[14] Several ECG indices thought to represent atrial remodeling have been independently associated with stroke and AF.^[15] These measures include the (i) PWD; (ii) PWDIs; (iii) PTFV₁ in the precordial lead V1; (iv) P-wave axis; and (v) interatrial blocks (IABs).^[16] Previous studies have identified several P-wave indices that are markers of LA dysfunction and are associated with ischemic stroke with or without AF.^[17] Previous studies have reported that maximum PWD may be used for the prediction of AF.^[18] However, we did not detect PWD to be a predictor for the presence of AF in our study.

P wave dispersion is considered to reflect impaired and heterogeneous interatrial conduction, which is a specific and sensitive marker of AF in a wide variety of conditions.^[19] Dilaveris et al.^[18] found that PWDIs was significantly higher in patients with paroxysmal AF compared to the control group, and a PWDIs value of 40 ms distinguished paroxysmal AF patients from the control group with a sensitivity of 83% and a specificity of 85%. Aytemir et al.^[20] reported PWDIs >36 ms to be an independent predictor for the development of AF with a sensitivity of 77% and specificity of 82%. The PWDIs has been used for the prediction of AF in several clinical situations such as hyperthyroidism, chronic obstructive pulmonary disease, acute ischemic stroke and hypertrophic cardiomyopathy.^[19] Doğan et al.^[15] reported PWDIs as an independent predictor for the development of AF in patients with acute ischemic stroke. Similarly, we also found PWDIs to be a predictor of PAF in patients with ischemic stroke, with a sensitivity of 71% and specificity of 69%.

The PTFV₁ was first used by Morris et al.^[21] in 1964 as a representative of LA overload in several valvular heart diseases. Later, PTFV₁ was found to be an indicator of various pathologies such

as increased LA pressure, LA hypertrophy, LA enlargement, and abnormal interatrial conduction.^[17] Since AF development is also associated with these structural changes and electrical remodeling, PTFV₁ may be a good predictor of AF development. PTFV₁ >4000 $\mu\text{V}\cdot\text{ms}$ is accepted to be abnormal. An abnormal PTFV₁ level has been shown to negatively affect prognosis in heart failure and myocardial infarction.^[22] It was reported that a 1-SD increase of PTFV₁ increased the risk of AF occurrence by 27%.^[17] Additionally, PTFV₁ was found to be a better predictor in hemodialysis and stroke patients compared to the normal population.^[17] The PTFV₁ is indicative of LA volume overload and it has, therefore, been frequently used for AF prediction in patients undergoing hemodialysis.^[17] Goda et al.^[23] found PTFV₁ to be a strong predictor of AF in patients with acute ischemic stroke. In addition, PTFV₁ was reported to be a good predictor of stroke, regardless of AF in a meta-analysis by He et al.^[24] However, Sajeev et al.^[25] suggested that PTFV₁ was a weak predictor of ischemic stroke. Similarly, we found that PTFV₁ had a lower sensitivity and specificity in the detection of AF compared to PWDIs.

There are some limitations in the current study. First, our sample size was relatively small, which may have weakened the strength of our results. Second, this study is retrospective in nature. Third, Holter ECG monitoring was performed for 24 to 48 h in all patients. If a longer follow-up could have been made, PAF attacks could have been detected in more patients.

In conclusion, PWDIs and PTFV₁ in lead V1 are independent predictors for the presence of PAF in patients with ischemic stroke. These simple and easily accessible predictors, which can be detected by surface ECG, may help in identifying patients that require longer rhythm monitoring to detect occult PAFs, thereby preventing recurrent strokes.

Declaration of conflicting interests

The authors declared no conflicts of interest with respect to the authorship and/or publication of this article.

Funding

The authors received no financial support for the research and/or authorship of this article.

REFERENCES

1. Hindricks G, Potpara T, Dagres N, Arbelo E, Bax JJ, Blomström-Lundqvist C, et al. 2020 ESC Guidelines for the

- diagnosis and management of atrial fibrillation developed in collaboration with the European Association for Cardio-Thoracic Surgery (EACTS): The Task Force for the diagnosis and management of atrial fibrillation of the European Society of Cardiology (ESC) Developed with the special contribution of the European Heart Rhythm Association (EHRA) of the ESC. *Eur Heart J* 2021;42:373-498.
2. Boriani G, Savelieva I, Dan GA, Deharo JC, Ferro C, Israel CW, et al. Chronic kidney disease in patients with cardiac rhythm disturbances or implantable electrical devices: Clinical significance and implications for decision making-a position paper of the European Heart Rhythm Association endorsed by the Heart Rhythm Society and the Asia Pacific Heart Rhythm Society. *Europace* 2015;17:1169-96.
 3. Smit MD, Van Gelder IC. New treatment options for atrial fibrillation: Towards patient tailored therapy. *Heart* 2011;97:1796-802.
 4. Wolf PA, Abbott RD, Kannel WB. Atrial fibrillation as an independent risk factor for stroke: The Framingham Study. *Stroke* 1991;22:983-8.
 5. Adams HP Jr, del Zoppo G, Alberts MJ, Bhatt DL, Brass L, Furlan A, et al. Guidelines for the early management of adults with ischemic stroke: A guideline from the American Heart Association/American Stroke Association Stroke Council, Clinical Cardiology Council, Cardiovascular Radiology and Intervention Council, and the Atherosclerotic Peripheral Vascular Disease and Quality of Care Outcomes in Research Interdisciplinary Working Groups: The American Academy of Neurology affirms the value of this guideline as an educational tool for neurologists. *Circulation* 2007;115:e478-534.
 6. Schnabel RB, Aspelund T, Li G, Sullivan LM, Suchy-Dicey A, Harris TB, et al. Validation of an atrial fibrillation risk algorithm in whites and African Americans. *Arch Intern Med* 2010;170:1909-17.
 7. Chamberlain AM, Agarwal SK, Folsom AR, Soliman EZ, Chambless LE, Crow R, et al. A clinical risk score for atrial fibrillation in a biracial prospective cohort (from the Atherosclerosis Risk in Communities [ARIC] study). *Am J Cardiol* 2011;107:85-91.
 8. European Heart Rhythm Association; European Association for Cardio-Thoracic Surgery, Camm AJ, Kirchhof P, Lip GY, Schotten U, Savelieva I, Ernst S, et al. Guidelines for the management of atrial fibrillation: The Task Force for the Management of Atrial Fibrillation of the European Society of Cardiology (ESC). *Eur Heart J* 2010;31:2369-429.
 9. Cardiogenic brain embolism. Cerebral Embolism Task Force. *Arch Neurol* 1986;43:71-84.
 10. Mahajan R, Perera T, Elliott AD, Twomey DJ, Kumar S, Munwar DA, et al. Subclinical device-detected atrial fibrillation and stroke risk: A systematic review and meta-analysis. *Eur Heart J* 2018;39:1407-15.
 11. Saver JL. Clinical Practice. Cryptogenic Stroke. *N Engl J Med* 2016;374:2065-74.
 12. Christophersen IE, Yin X, Larson MG, Lubitz SA, Magnani JW, McManus DD, et al. A comparison of the CHARGE-AF and the CHA2DS2-VASc risk scores for prediction of atrial fibrillation in the Framingham Heart Study. *Am Heart J* 2016;178:45-54.
 13. Emren SV, Kocabaş U, Duygu H, Levent F, Şimşek EÇ, Yapan Emren Z, et al. The role of HATCH score in predicting the success rate of sinus rhythm following electrical cardioversion of atrial fibrillation. *Kardiol Pol* 2016;74:978-84.
 14. Chen LY, Soliman EZ. P Wave indices-advancing our understanding of atrial fibrillation-related cardiovascular outcomes. *Front Cardiovasc Med* 2019;6:53.
 15. Dogan U, Dogan EA, Tekinalp M, Tokgoz OS, Aribas A, Akilli H, et al. P-wave dispersion for predicting paroxysmal atrial fibrillation in acute ischemic stroke. *Int J Med Sci* 2012;9:108-14.
 16. Li TYW, Yeo LLL, Ho JSY, Leow AS, Chan MY, Dalakoti M, et al. Association of electrocardiographic P-wave markers and atrial fibrillation in embolic stroke of undetermined source. *Cerebrovasc Dis* 2021;50:46-53.
 17. Huang Z, Zheng Z, Wu B, Tang L, Xie X, Dong R, et al. Predictive value of P wave terminal force in lead V1 for atrial fibrillation: A meta-analysis. *Ann Noninvasive Electrocardiol* 2020;25:e12739.
 18. Dilaveris PE, Gialafos EJ, Sideris SK, Theopistou AM, Andrikopoulos GK, Kyriakidis M, et al. Simple electrocardiographic markers for the prediction of paroxysmal idiopathic atrial fibrillation. *Am Heart J* 1998;135:733-8.
 19. Fujimoto Y, Yodogawa K, Takahashi K, Tsuboi I, Hayashi H, Uetake S, et al. Noninvasive evaluation of reverse atrial remodeling after catheter ablation of atrial fibrillation by P wave dispersion. *Heart Vessels* 2017;32:1375-81.
 20. Aytemir K, Ozer N, Atalar E, Sade E, Aksöyek S, Ovünç K, et al. P wave dispersion on 12-lead electrocardiography in patients with paroxysmal atrial fibrillation. *Pacing Clin Electrophysiol* 2000;23:1109-12.
 21. Morris JJ Jr, Estes EH Jr, Whalen RE, Thompson HK Jr, McIntosh HD. P-wave analysis in valvular heart disease. *Circulation* 1964;29:242-52.
 22. Li Q, Gu LD, Zhang C, Liu W, Peng Y, Chai H, et al. A predictive study of the dynamic development of the P-wave terminal force in lead V1 in the electrocardiogram in relation to long-term prognosis in non-ST-segment elevation acute coronary syndrome patients during hospitalization. *Ann Noninvasive Electrocardiol* 2015;20:542-53.
 23. Goda T, Sugiyama Y, Ohara N, Ikegami T, Watanabe K, Kobayashi J, et al. P-wave terminal force in lead V1 predicts paroxysmal atrial fibrillation in acute ischemic stroke. *J Stroke Cerebrovasc Dis* 2017;26:1912-5.
 24. He J, Tse G, Korantzopoulos P, Letsas KP, Ali-Hasan-Al-Saegh S, Kamel H, et al. P-wave indices and risk of ischemic stroke: A systematic review and meta-analysis. *Stroke* 2017;48:2066-72.
 25. Sajeev JK, Koshy AN, Dewey H, Kalman JM, Bhatia M, Roberts L, et al. Poor reliability of P-wave terminal force V1 in ischemic stroke. *J Electrocardiol* 2019;52:47-52.